Care pathways –Task 2

- Review (BMJ 2012 Sep 3;345:e5205) → Evidence
 on the care of patients with multimorbidity is
 limited... Interventions had mixed effects...
- Update (NIVEL) → Programs varied in the target patient groups, implementation settings, number of included interventions, and number of chronic care model components ... Different components of the intervention were identified (comprehensive programs)



Lack of standardization

Needs

- Identify components of interventions
- Assess their cost-effectiveness
- Identify target population for each component
- How to standardize components?
- Setting?

Needs

- Identify components of interventions
 - → based on the work by NIVEL

- Delivery system design
 - Case manager
 - Multidisciplinary team
 - Individualized care plans
 - Comprehensive assessment

- Decision support
 - Implementation of EBM
 - Team training

- Self management
 - Health litteracy
 - Patient education (verbal education + educational material)
 - Family education
 - Self-management approach for patients
 - Patient assessment of quality of care

- Community resources
 - Access to community resources
 - Involvement of social network
 - Psychosocial support

- Clinical information system
 - Electronic patients records
 - Computerized clinical charts

Needs

- Identify components of interventions
- Assess their cost-effectiveness
- Identify target population for each component
- How to standardize components?
- Setting

Expert meeting

- 1. Expert judgment
 - Effectiveness
 - Cost effectiveness
 - Reliability
 - Applicability
- 2. Identify target population for each component based on risk factors.
- 3. Define minimum set of key characteristics to standardize them

Your opinion

- Programme
- Components
- Expert meeting and issues to assess
- Experts?